

Treadmill running frequency on anxiety and hippocampal adenosine receptors density in adult and middle-aged rats

Marcelo S. Costa ^{*}, Ana Paula Ardais, Gabriela T. Fioreze, Sabrina Mioranza, Paulo Henrique S. Botton, Luis Valmor Portela, Diogo O. Souza, Lisiane O. Porciúncula

Laboratory of Studies on the Purinergic System, Graduation Program in Biological Sciences/Biochemistry, Federal University of Rio Grande do Sul, Health and Basic Sciences Institute, Department of Biochemistry, Porto Alegre/RS 90035-003, Brazil

ARTICLE INFO

Article history:

Received 1 August 2011

Received in revised form 20 October 2011

Accepted 20 October 2011

Available online 28 October 2011

Keywords:

Adenosine receptors

Aging

Anxiety

Physical exercise

Treadmill running

ABSTRACT

Physical exercise protocols have varied widely across studies raising the question of whether there is an optimal intensity, duration and frequency that would produce maximal benefits in attenuating symptoms related to anxiety disorders. Although physical exercise causes modifications in neurotransmission systems, the involvement of neuromodulators such as adenosine has not been investigated after chronic exercise training. Anxiety-related behavior was assessed in the elevated plus-maze in adult and middle-aged rats submitted to 8 weeks of treadmill running 1, 3 or 7 days/week. The speed of running was weekly adjusted to maintain moderate intensity. The hippocampal adenosine A₁ and A_{2A} receptors densities were also assessed. Treadmill running protocol was efficient in increasing physical exercise capacity in adult and middle-aged rats. All frequencies of treadmill running equally decreased the time spent in the open arms in adult animals. Middle-aged treadmill control rats presented lower time spent in the open arms than adult treadmill control rats. However, treadmill running one day/week reversed this age effect. Adenosine A₁ receptor was not changed between groups, but treadmill running counteracted the age-related increase in adenosine A_{2A} receptors. Although treadmill running, independent from frequency, triggered anxiety in adult rats and treadmill running one day/week reversed the age-related anxiety, no consistent relationship was found with hippocampal adenosine receptors densities. Thus, our data suggest that as a complementary therapy in the management of mental disturbances, the frequency and intensity of physical exercise should be taken into account according to age. Besides, this is the first study reporting the modulation of adenosine receptors after chronic physical exercise, which could be important to prevent neurological disorders associated to increase in adenosine A_{2A} receptors.

© 2011 Elsevier Inc. Open access under the [Elsevier OA license](http://creativecommons.org/licenses/by-nc-sa/3.0/).

1. Introduction

Over the past years, a wide range of studies have shown that increased physical activity or regular physical exercise promote positive effects on brain functions (Cotman and Berchtold, 2002; Dishman et al., 2006; Gillum and Obisesan, 2010; Kramer et al., 1999; Rolland et al., 2010). Otherwise, the sedentary lifestyle has been correlated with a higher risk for development of obesity, cardiovascular diseases, type 2 diabetes, osteoporosis, cancer and depression (Roberts and Barnard, 2005).

The prevalence of mental health disorders among the elderly is often unrecognized, but it has been estimated that one in four older adults presents symptoms of depression, anxiety disorders or other psychiatric disorders. Mental health disorders are frequently comorbid

in older adults, occurring with a number of common chronic diseases (Fiske et al., 2009; Flint, 2005). In this scenario, the prevalence of anxiety disorders is estimated to be as high as 25% (Craske et al., 2009; Hettema et al., 2001).

Adenosine is a neuromodulator in the central nervous system (CNS), which via mainly adenosine A₁ and A_{2A} metabotropic receptors control synaptic transmission of many neurotransmitters, such as glutamate and dopamine. Adenosine A₁ receptors are expressed throughout the brain while adenosine A_{2A} is more restricted to the basal ganglia (Cunha, 2001). The physical exercise influences the central dopaminergic, noradrenergic and serotonergic systems (Meeusen and De Meirleir, 1995). In a previous study, the participation of adenosine receptors in physical activity was reported, in which caffeine increased treadmill running time and spontaneous locomotor activity on the open field task, whereas the adenosine A₁/A_{2A} receptor agonist, 5'-N-ethylcarboxamido-adenosine (NECA) decreased it (Davis et al., 2003). However, the effects of chronic physical exercise on this important neuromodulator were not investigated yet.

Pharmacological approaches and genetic deletion of adenosine A₁ and A_{2A} receptors revealed that both receptors are implicated in the

Abbreviation: VO_{2max}, maximum oxygen uptake.

^{*} Corresponding author at: Federal University of Rio Grande do Sul, Department of Biochemistry, Rua Ramiro Barcelos, 2600, anexo, Porto Alegre/RS 90035-003, Brazil. Tel.: +55 51 3308 5557; fax: +55 51 3308 5540.

E-mail address: marsilcos@yahoo.com.br (M.S. Costa).

etiology of anxiety (for review see Cunha et al., 2008). The knockout mice for adenosine A₁ as well as for adenosine A_{2A} receptor display an exacerbation of anxiety-related behavior (Giménez-Llort et al., 2002; Johansson et al., 2001; Lang et al., 2003; Ledent et al., 1997). In addition, anxiogenic effects of caffeine, a recognized nonselective adenosine A₁ and A_{2A} receptor antagonist, are demonstrated in a wide range of rodent studies (File et al., 1988; Florio et al., 1998; Jain et al., 1995).

According to epidemiological studies, the physical activity can attenuate anxiety-related symptoms (Bhui and Fletcher, 2000; Dunn et al., 2001; Goodwin, 2003; Lampinen et al., 2000; Motl et al., 2004), being applied as a non-pharmacological intervention in patients, mainly in older adults (Kligman and Pepin, 1992). However, studies evaluating effects of exercise on anxiety-related behavior in rodents are more controversial, ranging from a reduction on anxiety (Greenwood et al., 2003), no effect (Pietropaolo et al., 2008), to even more anxiety after voluntary exercise (Burghardt et al., 2004; Leasure and Jones, 2008; Van Hoomissen et al., 2004).

Treadmill running is widely used by humans (Winter et al., 2007) because all variables implicated in physical training programs necessary to achieve great results can be controlled. Indeed, the frequency of the physical exercise is often controlled for maximizing the training and for avoiding negative effects of overtraining (ACSM, 1998). Nonetheless, effects of physical exercise on the anxiety-related behavior usually do not take into account the frequency of sessions and the intensity throughout training, which in turn can contribute for inconclusive results.

In this study, the effects of frequency of physical exercise were evaluated on anxiety-related behavior in adult and middle-aged rats for better understanding its effects on mental health. Considering the involvement of the adenosine receptors in anxiety disorders and its important role as a neuromodulator in the CNS, the density of adenosine A₁ and A_{2A} receptors was assessed in the hippocampus from adult and middle-aged rats trained on the treadmill running 1, 3 or 7 days/week during 8 weeks at moderate intensity. The hippocampus is a brain structure important in processes involving learning and memory, but its participation on anxiety has also been evidenced in other studies (Barkus et al., 2010; Oler et al., 2010; Sapolski et al., 1990; Xiang et al., 2011).

This schedule of exercise training (1, 3 and 7 days per week, 20 min per session at moderate intensity, for 8 weeks) was chosen to mimic exercise conditions in humans (ACSM, 1998; 2009). Thus, in this study we analyzed the effects of low and high exercise frequencies. Moreover, 8 weeks of training is considered sufficiently chronic to promote health-related adaptations.

2. Materials and methods

2.1. Animals

Adult (3 month-old) and middle-aged (14–16 month-old) male Wistar rats were used. The animals were housed into plastic cages under a light/dark cycle (lights on at 7:00 a.m.), with tap water and food ad libitum and kept at 23 °C. All procedures were carried out according to NIH Guide for Care and Use of Laboratory Animals, and Brazilian Society for Neuroscience and Behavior (SBNeC) recommendations for animal care. This work was approved by the ethical committee of Federal University of Rio Grande do Sul. All animals were killed by decapitation after behavioral analysis.

2.2. Treadmill exercise protocol

The treadmill exercise protocol was started from 8 to 10 a.m. Prior to the exercise training, animals were habituated in the treadmill apparatus for rat (INBRAMED TK 01, Porto Alegre, Brazil) during one week, in order to minimize novelty-induced stress. They were

randomly assigned to eight experimental groups according to age (adult and middle-aged rats): treadmill controls (0); treadmill running 1 day/week (1); treadmill running 3 days/week (3), and treadmill running 7 days/week (7).

At the first day of the second week, an incremental test in the treadmill was performed to indirectly determine the maximal oxygen uptake (VO₂max) as recommended by Brooks and White (1978). This test was to determine the running speed during the training period. Briefly, animals were placed in the treadmill to run during 25 min at a low initial speed. The speed was increased by 5 m/min each 3 min up to exhaustion of the animal. Exhaustion was achieved when animals could not respond to the increase of speed. The time to fatigue (in minutes) and workload (expressed by velocity in m/min) were taken as indexes of capacity for exercise, and as a measure of VO₂max. This measure was used to control the exercise intensity during the physical training program. The incremental test was repeated in the seventh week of treadmill running training to assess the efficacy of this protocol. The intensity of the physical training protocol was adapted for each animal and it never surpassed 60–75% of the respective maximum oxygen uptake. The exercise training protocol consisted of treadmill running during 8 weeks. Each training session lasted 20 min because great improvements in physical fitness were achieved at this time (ACSM, 1998). The velocity of treadmill was weekly increased for all treadmill groups. The treadmill controls (0) were always habituated to the experimental room and handled as treadmill runners but they were not submitted to the forced running protocol (adapted from Scopel et al., 2006). The treadmill controls were placed in the treadmill off 3 days/week during 20 min. Then, animals returned to their home cages. All groups were handled every day to avoid differences in handling in the elevated plus-maze. Importantly, in the frequency of 3 days/week the animals were not submitted to the treadmill running for 3 consecutive days. There was an interval of one day between exercise days. Three days/week for control groups was chosen because this was the intermediate frequency among groups.

The velocity of running at the beginning of training was 10 m/min and 7 m/min for adult rats and middle-aged rats, respectively. This velocity was weekly increased, and at the end of training reached 20 m/min and 12 m/min for adult and middle-aged rats, respectively. This procedure was essential to keep the intensity constant in 60–75% of the maximum oxygen uptake along the 8 weeks of treadmill running training.

2.3. Elevated plus-maze

The analysis in the elevated plus-maze took place during the early light portion of the light/dark cycle. All groups were exposed to the maze on Tuesday, 48 h after the last Sunday running session for all frequencies of exercise training. This time period between the last treadmill running session and analysis in the elevated plus-maze was chosen to avoid acute exercise-related stress (Contarteze et al., 2008), which could influence the effects of treadmill running training for 8 weeks. The apparatus was made of wood covered with a layer of black formica and had four elevated arms (52 cm from the floor) 50 cm long and 10 cm wide. The arms were arranged in a cross-like disposition, with two opposite arms being enclosed (by 40 cm high walls) and two being open, having at their intersection a central platform (10 × 13.5 cm) that gave access to any of the four arms. The open arms were surrounded by a raised ledge (1 mm thick and 5 mm high). The platform was under dim light (70 lx). Each rat was placed in the central platform facing an open arm and the following observations were manually recorded during 5 min by two observers blind to the experimental groups. Each animal was tested for 5 min. Four variables were measured: (1) time spent in the open arms; (2) number of entries into the open arms; (4) number of entries into the closed arms; and (4) number of rearing. The rat was considered to have entered or spent time in an arm only when all four paws were in the respective arm.

The values obtained for time and the number of entries in the open arms are taken as measures of anxiety reduction. In addition, the number of closed arms entered was taken as an index of general activity and the number of rearing as exploratory activity (Hogg, 1996; Pellow and File, 1986).

2.4. SDS-PAGE (sodium dodecyl sulfate-polyacrylamide) immunoblotting

Twenty-four hours after behavioral tests, rats were sacrificed by cervical displacement; the hippocampus was dissected out and immediately homogenized in 5% SDS with a protease inhibitor cocktail (Sigma, São Paulo/Brazil) and frozen at -70°C . A sample for each set of experiments was collected for western blotting analysis. After defrost, the protein content was determined by Bicinchoninic acid assay using bovine serum albumin (BSA) as standard (Pierce, São Paulo/Brazil). Samples extracts were diluted to a final protein concentration of $2\text{ }\mu\text{g}/\mu\text{L}$ in SDS-PAGE buffer. Forty micrograms of the samples and prestained molecular weight standards (Bio-Rad, São Paulo/Brazil) were separated by SDS-PAGE (12% with 4% concentrating gel). After electro-transfer, the membranes were blocked with Tris-buffered saline 0.1% Tween-20 (TBS-T) containing 3% BSA for 1 h, except to adenosine A_{2A} receptors, which was blocked for 20 min. The membranes were incubated overnight for rabbit anti-adenosine A_1 receptors (1:1000, Affinity Bioreagents, São Paulo/SP, Brazil) and goat anti-adenosine A_{2A} receptors (1:500, Santa Cruz Biotechnologies, São Paulo/SP, Brazil) at 4°C for 48 h and then re-probed for mouse anti- α -actin (1:3000, Santa Cruz Biotechnologies, São Paulo/SP, Brazil). After primary antibody incubation, membranes were washed and incubated with horseradish-peroxidase conjugated secondary antibodies for 2 h at room temperature and developed with chemiluminescence ECL kit (Amersham, São Paulo/Brazil). The autoradiographic films were scanned and densitometric analyses were performed using public domain NIH Image Program (developed at the U.S. National Institutes of Health and available on the internet at <http://rsb.info.nih.gov/ni-image/>). Results were measured in density units using actin as a control of the protein loading.

2.5. Statistical analysis

All data were expressed as mean \pm SEM and analyzed by two-way ANOVA (age \times frequency as factors), followed by Fisher PLSD test when appropriate. Repeated measures ANOVA were used to assess the effects of treadmill running training on $\text{VO}_{2\text{max}}$. Significance was established at $P < 0.05$. StatSoft Statistica 10.0 (StatSoft, Inc., USA) was the software used.

3. Results

The efficiency of our exercise training protocol was demonstrated by the performance in the second test to estimate $\text{VO}_{2\text{max}}$ in adult and middle-aged rats (Fig. 1). Treadmill running increased the performance in the second test in all trained rats, which was measured by the maximal speed throughout test [$F(1,113) = 4242.3$, $P < 0.00001$]. However, adult rats presented higher performance than middle-aged rats in all groups [$F(1,113) = 383.64$, $P < 0.00001$]. Moreover, treadmill running 3 and 7 days/week were more efficient to increase maximum oxygen uptake estimated in both age groups [$F(3,113) = 483.35$, $P < 0.00001$]. The effects of age and frequency of exercise in $\text{VO}_{2\text{max}}$ showed here were in accordance to human studies (ACSM, 1998). The second test was carried out in the seventh week and not in the end of the training to avoid the stress induced by test in the elevated plus-maze.

The time spent in the open arms in the elevated plus-maze was recorded for adult and middle-aged rats 48 h after the end of treadmill running training. Two-way ANOVA analysis revealed a significant interaction between frequency of running and age [$F(3,113) =$




Fig. 1. Effects of treadmill running protocol on physical exercise capacity in adult and middle-aged rats. Each bar represents mean \pm S.E.M of the time in the incremental test to estimated $\text{VO}_{2\text{max}}$ for 16–18 adult rats or 12–14 middle-aged per group; 0 – control; 1 – one day/week; 3 – three days/week and 7 – seven days/week. Rats ran in the treadmill during 8 weeks, 20 min each session. ** $P < 0.05$, indicates significant effects of age (0). * $P < 0.05$, indicates differences from their control counterparts. # $P < 0.05$ indicates difference from their treadmill running 1 day/week counterparts (1). The data were analyzed by two-way ANOVA followed by the Fisher PLSD test.

4.6889, $P = 0.004$] (Fig. 2). Adult treadmill control rats (0 – white bars) spent twofold more time in the open arms than middle-aged treadmill control rats (0 – gray bars) ($P = 0.0013$). For adult rats, all frequencies of treadmill running caused a similar decrease on the time spent in the open arms (45% of decrease) when compared to adult treadmill control rats [1 day/week $P = 0.002876$; 3 days/week $P = 0.004272$; 7 days/week $P = 0.006352$] (Fig. 2, white bars). For middle-aged rats, the frequency of one day/week increased the time spent in the open arms when compared to middle-aged treadmill control rats [$P = 0.0243311$] (Fig. 2, gray bars).

The closed arm entries in the elevated plus-maze were used as an index of the locomotor activity. Two-way ANOVA analysis revealed an effect of age [$F(1,113) = 5.6423$, $P = 0.0192$] (Fig. 3A). Two-way ANOVA analysis also revealed an effect of age in open arms entries [$F(1,113) = 4.4829$, $P = 0.03645$] (Fig. 3B). A significant interaction was observed in the number of entries in the open arms [$F(3,113) = 3.9729$, $P = 0.00986$]. Adult treadmill control rats (0 – white bars) present twofold more entries in the open arms than middle-aged treadmill control rats (0 – gray bars) ($P = 0.001815$). Besides, in adult rats, treadmill running one day/week was able to decrease the number of entries in the open arms when compared to non-runners [$P = 0.001236$].

As an index of exploratory activity, the number of rearing was recorded during 5 min of exploration in the elevated plus-maze. Two-way ANOVA analysis revealed only a significant effect of age [$F(1,113) = 24.227$, $P < 0.0001$]. Thus, middle-aged rats presented




Fig. 2. Time spent in the open arms in the elevated plus-maze for adult and middle-aged rats submitted to treadmill running training. Each bar represents mean \pm S.E.M of the time spent in the open arms (minutes) for 16–18 adult rats or 12–14 middle-aged per group; 0 – control; 1 – one day/week; 3 – three days/week and 7 – seven days/week. Rats ran in the treadmill during 8 weeks, 20 min each session. * $P < 0.05$, indicates difference from control adult rats (0). # $P < 0.05$, indicates differences from their control counterparts. The data were analyzed by two-way ANOVA followed by the Fisher PLSD test.




Fig. 3. Number of entries in the arms for adult and middle-aged rats in the elevated plus-maze submitted to treadmill running training. Each bar represents mean \pm S.E.M of (A) the number of entries in the closed arms and (B) the number of entries in the open arms for 16–18 adult rats or 12–14 middle-aged per group; 0 – control; 1 – one day/week; 3 – three days/week and 7 – seven days/week. Rats ran in the treadmill during 8 weeks, 20 min each session. * $P < 0.05$ indicates significant effects of age. # $P < 0.05$, indicates differences from treadmill running control adult (0). The data were analyzed by two-way ANOVA followed by the Fisher PLSD test.

lower number of rearing when compared to adults independent on the frequency of treadmill running (Fig. 4).

The immunocontent of whole hippocampal extracts was assessed for adenosine A_1 and A_{2A} receptors (Fig. 5A and B). No differences were found in the adenosine A_1 receptors (Fig. 5A). A significant effect of age was found for adenosine A_{2A} receptors density in the hippocampus [$F(1, 39) = 4.5979$; $P = 0.0383$]. In addition, the effect of frequency [$F(3, 39) = 8.408$; $P = 0.0002$] and a significant interaction was also observed [$F(3, 39) = 5.4817$; $P = 0.00306$]. Thus, all frequencies of treadmill running significantly prevented the age-related increase in the adenosine A_{2A} receptor density, with a robust decrease for frequency of 7 days/week [$P < 0.00001$] (Fig. 5B).

4. Discussion

In this study, 8 weeks of treadmill running was able to improve the physical exercise capacity, as demonstrated by the second test




Fig. 4. Number of rearing for adult and middle-aged rats submitted to treadmill running. Each bar represents mean \pm S.E.M of number of rearing for 16–18 adult rats or 12–14 middle-aged per group; 0 – control; 1 – one day/week; 3 – three days/week and 7 – seven days/week. Rats ran in the treadmill during 8 weeks, 20 min each session. * $P < 0.05$ indicates significant effects of age. The data were analyzed by two-way ANOVA followed by the Fisher PLSD test.




Fig. 5. Adenosine A_1 and A_{2A} receptors densities from the hippocampus of adult and middle-aged rats submitted to treadmill running. Data are expressed as means \pm SEM of density unit lines. (A) Adenosine A_1 receptors densities from adult ($n = 5-6$ per group) and middle-aged ($n = 6$ per group); (B) adenosine A_{2A} receptors densities from adult ($n = 5-7$ per group) and middle-aged ($n = 5-6$ per group); 0 – control; 1 – one day/week; 3 – three days/week and 7 – seven days/week. Rats ran in the treadmill during 8 weeks, 20 min each session. * $P < 0.05$, indicates significant effects of age (0). ** $P < 0.05$, indicates difference from control middle-aged rats (0) *** $P < 0.05$, indicates significant difference from all frequencies in middle-aged groups. The data were analyzed by two-way ANOVA followed by the Fisher PLSD test.

used to estimate VO_{2max} in adult and middle-aged rats. Although adult rats presented higher performance than middle-aged rats in all groups, the treadmill running increased the performance in the second test in all trained rats. The frequencies of treadmill running 3 and 7 days/week presented a more robust effect on the estimated maximum oxygen uptake for both ages. The effects of age and frequency of exercise in VO_{2max} were similar to that found in human studies (ACSM, 1998).

In the elevated plus-maze performance, distinct effects in adult and middle-aged rats could be observed. Initially, age differences could be confirmed independent on the frequency of treadmill running. Indeed, our findings confirmed that middle-aged rats commonly exhibit a more exacerbated anxiety-like behavior than adult rats (Boguszewski and Zagrodzka, 2002; Frussa-Filho et al., 1992; Moretti et al., 2011). Besides, middle-aged rats usually present a decrease in the locomotor and exploratory activities when compared to adult ones (Agafonova et al., 2007; Stefanova et al., 2010). In this context, middle-aged animals presented less number of entries in the closed as well as in the open arms and in the number of rearing compared to adults.

The frequency of exercise preserved the habitual locomotor and exploratory activities of the animals from both ages. The frequency only modified the time spent and the number of entries in the open arms. Even though differences found in the number of entries in the closed arms (as an index of locomotor activity) can limit the interpretation of the behavior in the elevated plus maze (Lister, 1990; Pellow et al., 1985), the effects of frequency seem to be related to the

anxiety-like behavior rather than locomotion for some reasons. Initially, number of entries in the arms is taken as a measure of locomotion, but current multifactorial studies have shown that this measure loads on two factors (anxiety and locomotion), whereas the number of entries in the closed arms loads exclusively (and with the highest loading) on the locomotion factor (File, 1991; Rodgers and Johnson, 1995). Since treadmill running did not affect the number of entries in the closed arms, the results found for adult rats suggest that all frequencies enhanced anxiety-like behavior. The frequency of one day/week attenuated anxiety-like behavior usually displayed by middle-aged animals.

The exercise abstinence of 48 h could explain these findings in adult rats, because anxiety is very common in trained humans during extended period of exercise abstinence (reviewed by Peluso and Guerra de Andrade, 2005). Nevertheless, this interval is necessary in order to avoid effects of fatigue caused by acute exercise as well as alterations in the locomotor activity by short intervals between the end of training and behavioral testing (Burghardt et al., 2004; Contarteze et al., 2008; Duman et al., 2008).

The impact of the physical exercise on anxiety-related behavior has reported contradictory findings in rodents. These discrepancies may be due to some factors, which include the type of exercise protocols (forced or voluntary), duration, intensity, frequency and behavioral paradigms used to assess physical exercise outcomes. Even though voluntary wheel running is largely used in studies with rodents, types of exercise will not be discussed here because this form of physical exercise does not accurately control intensity, duration and frequency of running. Furthermore, our protocol aimed to evaluate variables in the treadmill running because this type of exercise is widely used by humans.

Previous studies with behavioral analysis in the elevated plus-maze revealed that treadmill running 5 days a week during 10 weeks promoted an anxiolytic effect in adult rats (Fulk et al., 2004), whereas no effect was observed after 8 weeks of treadmill running, 5 days a week during 45 min at 20 m/min (Burghardt et al., 2004). In a recent study, adult rats that ran daily in a treadmill during 4 weeks presented normal performance in the light–dark box (Salim et al., 2010). The duration of training in these studies was similar, being considered chronic, and the interval between the end of training and the beginning of behavioral analysis was 48 h in order to avoid exercise-induced fatigue and stress (Duman et al., 2008). However, differently from other aforementioned studies in adult rats (Burghardt et al., 2004; Fulk et al., 2004), the speed of running in our protocol was weekly adjusted to maintain the intensity at 60–75% VO_2max according to the improvement of the physical capacity (see Fig. 1). It is important to emphasize that this control is crucial to maintain the intensity of running constant and promote improvement on the physical fitness (ACSM, 1998).

Our protocol revealed that only the frequency of one day a week of treadmill running was able to attenuate the anxiety-related behavior in middle-aged rats. Although the physical exercise does not interrupt the aging process, there is evidence that regular exercise can minimize detrimental effects of sedentary lifestyle and increase life expectancy by limiting the development and progression of chronic diseases and disabling conditions (ACSM, 2009; Harvey et al., 2010; Lazarov et al., 2010). The aging exacerbates anxiety-related behavior (File, 1990), and physical exercise may attenuate it (Cassilhas et al., 2010).

In this study, the protocol applied to the animals aimed to control the variables that affect outcomes of physical exercise, such as intensity, duration and frequency among training period. These variables are poorly controlled in other protocols of physical exercise used in rodents, which difficult interpretations of the findings about effects of exercise on anxiety. Moreover, this control is crucial to mimic physical exercise conditions applied in training protocols for human subjects. Here, eight weeks of moderate treadmill running for adult

rats decreased the time spent in open arms, independently of exercise frequency. However, treadmill running once a week reversed the age-related anxiety. Thus, this study clearly demonstrated that the frequency of exercise affects anxiety-related behavior depending on age and reveals the importance of the amount of physical exercise necessary to promote health and fitness. The period of training (8 weeks) is considered chronic, simulates training conditions in humans and promotes health-related adaptations. The responses from acute exercise are much attributed to an increase in the levels of stress-related measurements (Winter et al., 2007).

Over the past years, the impact of different types of physical exercise on many neurotransmission systems has been investigated in rodents, and some modifications in the glutamatergic, dopaminergic and GABAergic systems were found in different brain areas (Dietrich et al., 2005; Hill et al., 2010; Real et al., 2010; Sutoo and Akiyama, 2003). The participation of adenosine receptors in physical activity was reported by administration of caffeine and the adenosine $\text{A}_1/\text{A}_{2\text{A}}$ receptor agonist, 5'-N-ethylcarboxamidoadenosine (NECA) in rats (Davis et al., 2003). In this study, caffeine increased the running time in the treadmill and the spontaneous locomotor activity in the open field whereas the activation of $\text{A}_1/\text{A}_{2\text{A}}$ receptor had opposite effects. Besides, caffeine was able to prevent fatigue induced by NECA in the treadmill running. It is important to consider that in our study treadmill running did not affect exploration and locomotion in the elevated plus-maze. Thus, an acute manipulation of adenosine receptors seems to influence the physical exercise performance. In our study, modifications on adenosine $\text{A}_{2\text{A}}$ receptor density by chronic physical exercise were found. In fact, the hippocampal adenosine A_1 receptors were neither modified by frequencies of treadmill running nor by aging. Then, adenosine A_1 receptors seem not to be involved in the effects promoted by chronic treadmill running in adult and middle-aged rats. It could not be discarded that modifications might be occurring in this receptor by acute exercise.

In agreement with previous reports, the density of $\text{A}_{2\text{A}}$ receptors increased in the hippocampus from middle-aged rats when compared to adult animals (Canas et al., 2009; Castillo et al., 2009). The most direct evidence implicating adenosine receptors in anxiety comes from studies in rodents and humans reporting the anxiogenic effects of caffeine (for review see Cunha et al., 2008; El Yacoubi et al., 2000). Currently, the anxiogenic effect of caffeine in psychiatric patients and in healthy subjects may be related to the presence of polymorphisms in the adenosine $\text{A}_{2\text{A}}$ receptor gene (Alsene et al., 2003; Childs et al., 2008; Rogers et al., 2010; Tsai et al., 2006). Although all frequencies of treadmill running reversed the age-related increase on density of adenosine $\text{A}_{2\text{A}}$ receptors, the same could not be observed for anxiety-related behavior. In fact, only the frequency of one day a week presented modifications on adenosine $\text{A}_{2\text{A}}$ receptors along with attenuation in the anxiety-related behavior by aging. Even though modifications in the performance of plus maze have not been observed for frequencies of 3 and 7 days a week in middle-aged animals, physical exercise was able to prevent modifications in the adenosine $\text{A}_{2\text{A}}$ receptors by aging. Considering that increases in the density and expression of adenosine $\text{A}_{2\text{A}}$ receptors are associated with neurological disorders related to aging (Castillo et al., 2009; Cunha et al., 1995; Lopes et al., 1999; Meyer et al., 2007; Pagonopoulou and Angelatou, 1992), it remains to be determined whether other beneficial effects of physical exercise could be related to adenosine $\text{A}_{2\text{A}}$ receptors.

The alterations in neuroendocrine tissues and immune responses are associated with chronic stress and similar alterations have been shown by treadmill running in adult rats (Moraska et al., 2000). The protocol applied in this study suggests that the adequate control of the intensity and frequency is indeed crucial to ensure better results and to avoid health negative effects. Besides, our data suggest that middle-aged rats had benefits of moderate exercise but only at low frequency.

Physical exercise is considered a complementary therapy in the management of mild-to-moderate mental disturbances, which include depression and anxiety (Paluska and Schwenk, 2000). Our findings contribute for the importance in controlling all variables that could generate psychological symptoms deleterious to health (Peluso and Guerra de Andrade, 2005).

Acknowledgments

The authors are grateful for Brazilian Funding Agencies: PRONEX and PRONEX/FAPERGS, CNPq (Proc no. 472216/2009-9, Lisiane O. Porciúncula), CNPq/INCT – Excitotoxicity and Neuroprotection, and CAPES for the fellowship. Gabriela T. Fioreze is a recipient from CNPq fellowship.

References

- ACSM. Position stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc* 1998;30:975–91.
- ACSM. Position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc* 2009;41:1510–30.
- Agafonova IG, Kolosova NG, Mishchenko NP, Chaikina EL, Stonik VA. Effect of histochrome on brain vessels and research and exploratory activity of senescence accelerated OXYS rats. *Bull Exp Biol Med* 2007;143:467–71.
- Alsene K, Deckert J, Sand P, de Wit H. Association between A2a receptor gene polymorphisms and caffeine-induced anxiety. *Neuropsychopharmacology* 2003;28:1694–702.
- Barkus C, McHugh SB, Sprengel R, Seeburg PH, Rawlins JNP, Bannerman DM. Hippocampal NMDA receptors and anxiety: at the interface between cognition and emotion. *Eur J Pharmacol* 2010;626:49–56.
- Bhui K, Fletcher A. Common mood and anxiety states: gender differences in the protective effect of physical activity. *Soc Psychiatry Psychiatr Epidemiol* 2000;35:28–35.
- Boguszewski P, Zagrodzka J. Emotional changes related to age in rats: a behavioral analysis. *Behav Brain Res* 2002;133:323–32.
- Brooks GA, White TP. Determination of metabolic and heart rate responses of rats to treadmill exercise. *J Appl Physiol* 1978;45:1009–15.
- Burghardt PR, Fulk LJ, Hand GA, Wilson MA. The effects of chronic treadmill and wheel running on behavior in rats. *Brain Res* 2004;1019:84–96.
- Canas PM, Duarte JM, Rodrigues RJ, Köfalvi A, Cunha RA. Modification upon aging of the density of presynaptic modulation systems in the hippocampus. *Neurobiol Aging* 2009;30:1877–84.
- Cassilhas RC, Antunes HK, Tufik S, de Mello MT. Mood, anxiety, and serum IGF-1 in elderly men given 24 weeks of high resistance exercise. *Percept Mot Skills* 2010;110:265–76.
- Castillo CA, Albasan JL, Leon D, Jordan J, Pallas M, Camins A, et al. Age-related expression of adenosine receptors in brain from the senescence-accelerated mouse. *Exp Gerontol* 2009;44:453–61.
- Childs E, Hohoff C, Deckert J, Xu K, Badner J, de Wit H. Association between ADORA2A and DRD2 polymorphisms and caffeine-induced anxiety. *Neuropsychopharmacology* 2008;33:2791–800.
- Contarteze RV, Manchado FB, Gobatto CA, De Mello MA. Stress biomarkers in rats submitted to swimming and treadmill running exercises. *Comp Biochem Physiol A Mol Integr Physiol* 2008;151:415–22.
- Cotman CW, Berchtold NC. Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci* 2002;25:295–301.
- Craske MG, Rauch SL, Ursano R, Prenoveau J, Pine DS, Zinbarg RE. What is an anxiety disorder? *Depress Anxiety* 2009;26:1066–85.
- Cunha RA. Adenosine as a neuromodulator and as a homeostatic regulator in the nervous system: different roles, different sources and different receptors. *Neurochem Int* 2001;38:107–25.
- Cunha RA, Constantino MC, Sebastião AM, Ribeiro JA. Modification of A1 and A2a adenosine receptor binding in aged striatum, hippocampus and cortex of the rat. *Neuroreport* 1995;6:1583–8.
- Cunha RA, Ferré S, Vaugeois JM, Chen JF. Potential therapeutic interest of adenosine A2A receptors in psychiatric disorders. *Curr Pharm Des* 2008;14:1512–24.
- Davis JM, Zhao Z, Stock HS, Mehl KA, Buggy J, Hand GA. Central nervous system effects of caffeine and adenosine on fatigue. *Am J Physiol Regul Integr Comp Physiol* 2003;284:399–404.
- Dietrich MO, Mantese CE, Porciuncula LO, Ghisleni G, Vinade L, Souza DO, et al. Exercise affects glutamate receptors in postsynaptic densities from cortical mice brain. *Brain Res* 2005;1065:20–5.
- Dishman RK, Berthoud HR, Booth FW, Cotman CW, Edgerton VR, Fleshner MR, et al. Neurobiology of exercise. *Obesity* 2006;14:345–56.
- Duman CH, Schlesinger L, Russell DS, Duman RS. Voluntary exercise produces antidepressant and anxiolytic behavioral effects in mice. *Brain Res* 2008;1199:148–58.
- Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose response effects on outcome of depression and anxiety. *Med Sci Sports Exerc* 2001;33:S587–97.
- El Yacoubi M, Ledent C, Parmentier M, Costentin J, Vaugeois JM. The anxiogenic-like effect of caffeine in two experimental procedures measuring anxiety in the mouse is not shared by selective A(2A) adenosine receptor antagonists. *Psychopharmacology* 2000;148:153–63.
- File SE. New strategies in the search for anxiolytics. *Drug Des Deliv* 1990;5:195–201.
- File SE. The biological basis of anxiety. In: Meltzer HY, Nerozzi D, editors. *Current practices and future developments in the pharmacotherapy of mental disorders*. Amsterdam: Elsevier; 1991. p. 159–65.
- File SE, Baldwin HA, Johnston AL, Wilks LJ. Behavioral effects of acute and chronic administration of caffeine in the rat. *Pharmacol Biochem Behav* 1988;30:809–15.
- Fiske A, Wetherell JL, Gatz M. Depression in older adults. *Annu Rev Clin Psychol* 2009;5:363–89.
- Flint AJ. Generalised anxiety disorder in elderly patients: epidemiology, diagnosis and treatment options. *Drugs Aging* 2005;22:101–14.
- Florio C, Prezioso A, Papaioannou A, Vertua R. Adenosine A1 receptors modulate anxiety in CD1 mice. *Psychopharmacology* 1998;136:311–9.
- Frussa-Filho R, Otoboni JR, Giannotti AD, Amaral AC, Conceição IM. Effect of age on antinociceptive effects of elevated plus-maze exposure. *Braz J Med Biol Res* 1992;25:827–9.
- Fulk LJ, Stock HS, Lynn A, Marshall J, Wilson MA, Hand GA. Chronic physical exercise reduces anxiety-like behavior in rats. *Int J Sports Med* 2004;25:78–82.
- Gillum RF, Obisesan TO. Physical activity, cognitive function, and mortality in a US national cohort. *Ann Epidemiol* 2010;20:251–7.
- Giménez-Llort L, Fernández-Teruel A, Escorihuela RM, Fredholm BB, Tobeña A, Pekny M, et al. Mice lacking the adenosine A1 receptor are anxious and aggressive, but are normal learners with reduced muscle strength and survival rate. *Eur J Neurosci* 2002;16:547–50.
- Goodwin RD. Association between physical activity and mental disorders among adults in the United States. *Prev Med* 2003;36:698–703.
- Greenwood BN, Foley TE, Day HE, Campisi J, Hammack SH, Campeau S, et al. Freewheel running prevents learned helplessness/behavioral depression: role of dorsal raphe serotonergic neurons. *J Neurosci* 2003;23:2889–98.
- Harvey SB, Hotopf M, Overland S, Mykletun A. Physical activity and common mental disorders. *Br J Psychiatry* 2010;197:357–64.
- Hettema JM, Neale MC, Kendler KS. A review and meta-analysis of the genetic epidemiology of anxiety disorders. *Am J Psychiatry* 2001;158:1568–78.
- Hill LE, Droste SK, Nutt DJ, Linthorst AC, Reul JM. Voluntary exercise alters GABA(A) receptor subunit and glutamic acid decarboxylase-67 gene expression in the rat forebrain. *J Psychopharmacol* 2010;24:745–56.
- Hogg S. A review of the validity and variability of the elevated plus-maze as an animal model of anxiety. *Pharmacol Biochem Behav* 1996;54:21–30.
- Jain N, Kemp N, Adeyemo O, Buchanan P, Stone TW. Anxiolytic activity of adenosine receptor activation in mice. *Br J Pharmacol* 1995;116:2127–33.
- Johansson B, Halldner L, Dunwiddie TV, Masino SA, Poelchen W, Giménez-Llort L, et al. Hyperalgesia, anxiety, and decreased hypoxic neuroprotection in mice lacking the adenosine A1 receptor. *Proc Natl Acad Sci U S A* 2001;98:9407–12.
- Kligman EW, Pepin E. Prescribing physical activity for older patients. *Geriatrics* 1992;47:33–4. [37–44, 47].
- Kramer AF, Hahn S, Cohen NJ, Banich MT, McAuley E, Harrison CR, et al. Ageing, fitness and neurocognitive function. *Nature* 1999;400:418–9.
- Lampinen P, Heikkinen RL, Ruoppila I. Changes in intensity of physical exercise as predictors of depressive symptoms among older adults: an eight year follow-up study. *Prev Med* 2000;12:113–80.
- Lang UE, Lang F, Richter K, Vallon V, Lipp HP, Schnermann J, et al. Emotional instability but intact spatial cognition in adenosine receptor 1 knock out mice. *Behav Brain Res* 2003;145:179–88.
- Lazarov O, Mattson MP, Peterson DA, Pimplikar SW, van Praag H. When neurogenesis encounters aging and disease. *Trends Neurosci* 2010;33:569–79.
- Leasure JL, Jones M. Forced and voluntary exercise differentially affect brain and behavior. *Neuroscience* 2008;156:456–65.
- Ledent C, Vaugeois JM, Schiffmann SN, Pedrazzini T, El Yacoubi M, Vanderhaeghen JJ, et al. Aggressiveness, hypoalgesia and high blood pressure in mice lacking the adenosine A2a receptor. *Nature* 1997;388:674–8.
- Lister RG. Ethologically-based animal models of anxiety disorders. *Pharmacol Ther* 1990;46:321–40.
- Lopes LV, Cunha RA, Ribeiro JA. Cross talk between A1 and A2a adenosine receptors in the hippocampus and cortex of young adult and old rats. *J Neurophysiol* 1999;82:3196–203.
- Meeusen R, De Meirleir K. Exercise and brain neurotransmission. *Sports Med* 1995;20:160–88.
- Meyer PT, Elmenhorst D, Boy C, Winz O, Matusch A, Zilles K, et al. Effect of aging on cerebral A1 adenosine receptors: a [18F]CPFPX PET study in humans. *Neurobiol Aging* 2007;28:1914–24.
- Moraska A, Deak T, Spencer RL, Roth D, Fleshner M. Treadmill running produces both positive and negative physiological adaptations in Sprague–Dawley rats. *Am J Physiol Regul Integr Comp Physiol* 2000;279:R1321–9.
- Moretti M, de Souza AG, de Chaves G, de Andrade VM, Romão PR, Gavioli EC, et al. Emotional behavior in middle-aged rats: implications for geriatric psychopathologies. *Physiol Behav* 2011;102:115–20.
- Motl RW, Birnbaum AS, Kubik MY, Dishman RK. Naturally occurring changes in physical activity are inversely related to depressive symptoms during early adolescence. *Psychosom Med* 2004;66:336–42.
- Oler JA, Fox AS, Shelton SE, Rogers J, Dyer TD, Davidson RJ, et al. Amygdalar and hippocampal substrates of anxious temperament differ in their heritability. *Nature* 2010;466:864–8.
- Pagonopoulou O, Angelatou F. Reduction of A1 adenosine receptors in cortex, hippocampus and cerebellum in ageing mouse brain. *Neuroreport* 1992;3:735–77.
- Paluska SA, Schwenk TL. Physical activity and mental health: current concepts. *Sports Med* 2000;29:167–80.
- Pellow S, File SE. Anxiolytic and anxiogenic drug effects on exploratory activity in an elevated plus-maze: a novel test of anxiety in the rat. *Pharmacol Biochem Behav* 1986;24:525–9.

- Pellow S, Chopin P, File SE, Briley M. Validation of open: closed arm entries in an elevated plus-maze as a measure of anxiety in the rat. *J Neurosci Methods* 1985;14:149–67.
- Peluso MA, Guerra de Andrade LH. Physical activity and mental health: the association between exercise and mood. *Clinics* 2005;60:61–70.
- Pietropaolo S, Sun Y, Li R, Brana C, Feldon J, Yee BK. The impact of voluntary exercise on mental health in rodents: a neuroplasticity perspective. *Behav Brain Res* 2008;192:42–60.
- Real CC, Ferreira AF, Hernandez MS, Britto LR, Pires RS. Exercise-induced plasticity of AMPA-type glutamate receptor subunits in the rat brain. *Brain Res* 2010;1363:63–71.
- Roberts CK, Barnard RJ. Effects of exercise and diet on chronic disease. *J Appl Physiol* 2005;98:3–30.
- Rodgers RJ, Johnson JN. Factor analysis of spatiotemporal and ethological measures in the murine elevated plus-maze test of anxiety. *Pharmacol Biochem Behav* 1995;52:297–303.
- Rogers PJ, Hohoff C, Heatherley SV, Mullings EL, Maxfield PJ, Evershed RP, et al. Association of the anxiogenic and alerting effects of caffeine with ADORA2A and ADORA1 polymorphisms and habitual level of caffeine consumption. *Neuropsychopharmacology* 2010;35:1973–83.
- Rolland Y, Abellan Van Kan G, Vellas B. Healthy brain aging: role of exercise and physical activity. *Clin Geriatr Med* 2010;26:75–87.
- Salim S, Sarraj N, Taneja M, Saha K, Tejada-Simon MV, Chugh G. Moderate treadmill exercise prevents oxidative stress-induced anxiety-like behavior in rats. *Behav Brain Res* 2010;208:545–52.
- Sapolski RM, Uno H, Rebert CS, Finch CE. Hippocampal damage associated with prolonged glucocorticoid exposure in primates. *J Neurosci* 1990;10:2897–902.
- Scopel D, Fochesatto C, Cimarosti H, Rabbo M, Belló-Klein A, Salbego C, et al. Exercise intensity influences cell injury in rat hippocampal slices exposed to oxygen and glucose deprivation. *Brain Res Bull* 2006;71:155–9.
- Stefanova NA, Fursova AZh, Kolosova NG. Behavioral effects induced by mitochondria-targeted antioxidant SkQ1 in Wistar and senescence-accelerated OXYS rats. *J Alzheimers Dis* 2010;21:479–91.
- Sutoo D, Akiyama K. Regulation of brain function by exercise. *Neurobiol Dis* 2003;13:1–14.
- Tsai SJ, Hong CJ, Hou SJ, Yen FC. Association study of adenosine A2a receptor (1976C>T) genetic polymorphism and mood disorders and age of onset. *Psychiatr Genet* 2006;16:185.
- Van Hoomissen JD, Holmes PV, Zellner AS, Poudevigne A, Dishman RK. Effects of beta-adrenoreceptor blockade during chronic exercise on contextual fear conditioning and mRNA for galanin and brain-derived neurotrophic factor. *Behav Neurosci* 2004;118:1378–90.
- Winter B, Breitenstein C, Mooren FC, Voelker K, Fobker M, Lechtermann A, et al. High impact running improves learning. *Neurobiol Learn Mem* 2007;87:597–609.
- Xiang X, Huang W, Haile CN, Kosten TA. Hippocampal GluR1 associates with behavior in the elevated plus maze and shows sex differences. *Behav Brain Res* 2011;222:326–31.